

VACCINE SCHEDULE

<u>At birth:</u>	Hepatitis B (<i>given in the hospital</i>)
<u>2 months:</u>	Pentacel (DTaP-Hib-Polio), Prevnar, Hepatitis B, & Rotateq (<i>oral</i>)
<u>4 months:</u>*	Pentacel (DTaP-Hib-Polio), Prevnar, & Rotateq (<i>oral</i>)
<u>6 months:</u>**	Pentacel (DTaP-Hib-Polio), Prevnar, & Rotateq (<i>oral</i>)
<u>9 months:</u>	Hepatitis B
<u>12 months:</u>	Proquad (MMR-Chicken Pox), Prevnar, & Hepatitis A
<u>15 months:</u>	DTaP-Hib
<u>18 months:</u>	Hepatitis A (<i>must be a minimum of 6 months after the 1st Hepatitis A dose</i>)
<u>4 years:</u>	DTaP, Polio, & ProQuad (<i>must be after the 4th birthday</i>)
<u>11 years:</u>	Adacel & Menactra (<i>must be after the 11th birthday</i>)
<u>12 & up:</u>	Gardasil***
<u>16-17 yrs:</u>	Menactra & Bexsero

* If the first dose of Hepatitis B is NOT given at birth in the hospital, then a second dose will be given at age 4 months.

** Flu vaccine is recommended annually starting at age 6 months old. The first year a child receives Flu vaccine they will require a one-time booster one month after the first dose. Flumist (nasal spray) can be given to ages 2 years and up.

*** If the first dose of Gardasil is given BEFORE their 15th birthday, then your child only needs a one-time second dose (booster) 6 months later. If the first dose of Gardasil is given on or after their 15th birthday, then your child will require two additional doses (boosters): one booster 2 months after the first dose, and another booster would be four months after the second dose.

**** *This schedule is according to the American Academy of Pediatrics guidelines. It may change at any time. In addition, new vaccines may be introduced. If there are any changes, you will be informed.*